VOL. 459 JOURNAL OF CHROMATOGRAPHY DECEMBER 28, 1988

CONTENTS

12th INTERNATIONAL SYMPOSIUM ON COLUMN LIQUID CHROMATOGRAPHY, WASHINGTON, DC, JUNE 19–24, 1988, PART II

Obituary: István Halász (1922–1988)
by Cs. Horváth XIII

Design of optimized high-performance liquid chromatographic gradients for the separation of either small or large molecules. I. Minimizing errors in computer simulations
by B. F. D. Ghrist (Wilmington, DE, U.S.A.), B. S. Cooperman (Philadelphia, PA, U.S.A.) and L. R. Snyder (Orinda, CA, U.S.A.) 1

Design of optimized high-performance liquid chromatographic gradients for the separation of either small or large molecules. II. Background and theory
by B. F. D. Ghrist (Wilmington, DE, U.S.A.) and L. R. Snyder (Orinda, CA, U.S.A.) . . 25

Design of optimized high-performance liquid chromatographic gradients for the separation of either small or large molecules. III. An overall strategy and its application to several examples
by B. F. D. Ghrist (Wilmington, DE, U.S.A.) and L. R. Snyder (Orinda, CA, U.S.A.) . . 43

Effects of peak tailing on computer optimisation procedures for high-performance liquid chromatography. I. Characteristics of tailed peaks under optimisation conditions
by S. Sekulic and P. R. Haddad (Kensington, Australia) 65

Effects of peak tailing on computer optimisation procedures for high-performance liquid chromatography. II. An optimisation routine for tailed peaks
by P. R. Haddad and S. Sekulic (Kensington, Australia) 79

Universal detection and quantitation of surfactants by high-performance liquid chromatography by means of the evaporative light-scattering detector
by G. R. Bear (Bellaire, TX, U.S.A.) 91

Enhanced performance of a laser-induced fluorescence liquid chromatographic apparatus: a systems approach
by T. J. Edkins and D. C. Shelly (Hoboken, NJ, U.S.A.) 109

Experimental and theoretical model of refractive index artifacts in absorbance detection
by C. E. Evans, J. G. Shabushnig and V. L. McGuffin (East Lansing, MI, U.S.A.) . . . 119

Rhodamine labelling reagent for the determination of chlorophenols by liquid chromatography with peroxyoxalate chemiluminescence detection
by P. J. M. Kwakman, J. G. J. Mol, D. A. Kamminga, R. W. Frei, U. A. Th. Brinkman and G. J. de Jong (Amsterdam, The Netherlands) 139

Oxidative detection of coulometrically reduced organonitro pesticides in reversed-phase high-performance liquid chromatography
by R. T. Krause and Y. Wang (Washington, DC, U.S.A.) 151

Carbon–polymer chips as sensitive electrochemical detectors for micro-liquid chromatography
by L. J. Nagels (Antwerp, Belgium), J. M. Kauffmann (Brussels, Belgium), G. Schuddinck and C. Dewaele (Ghent, Belgium), G. J. Patriarche (Brussels, Belgium) and M. Verzele (Ghent, Belgium) 163

Water as a stationary phase modifier in packed-column supercritical fluid chromatography. I. Separation of free fatty acids
by F. O. Geiser (Glen Mills, PA, U.S.A.) and S. G. Yocklovich, S. M. Lurcott, J. W. Guthrie and E. J. Levy (Avondale, PA, U.S.A.) 173

Effect of the partial molar volume of the solute in the stationary phase on retention in supercritical fluid chromatography
by C. R. Yonker and R. D. Smith (Richland, WA, U.S.A.) 183

Use of two simultaneous detectors in capillary supercritical fluid chromatography
by D. J. Bornhop, S. Schmidt and N. L. Porter (Salt Lake City, UT, U.S.A.) 193

Effect of sample size on retention in packed column supercritical fluid chromatography. A method for characterizing stationary phase homogeneity
by P. J. Schoenmakers, L. G. M. Uunk and P. K. de Bokx (Eindhoven, The Netherlands) 201

Determination of a new inotropic agent in human plasma by high-performance liquid chromatography
by S.-H. Hsu, T. L. Koerper, B. J. Tomlinson, J. R. Miksic and P. E. Grebow (Horsham, PA, U.S.A.) 215

Separation of fatty acid binding protein by high-performance mixed-mode chromatography
by A. Samanta, G. A. Cordis, M. R. Prasad and D. K. Das (Farmington, CT, U.S.A.) . . 221

Novel dual-wavelength monitoring approach for the improved rapid separation and estimation of adenine nucleotides and creatine phosphate by high-performance liquid chromatography
by G. A. Cordis (Farmington, CT, U.S.A.), R. M. Engelman (Springfield, MA, U.S.A.) and D. K. Das (Farmington, CT, U.S.A.) 229

Quantitation of free amino acids in biological samples by high-performance liquid chromatography. Application of the method in evaluating amino acid levels in cerebrospinal fluid and plasma of patients with multiple sclerosis
by G. A. Qureshi and M. S. Baig (Huddinge, Sweden) 237

Determination of salsolinol by ion-exchange chromatography with glycylglycine as the post-derivatizing agent
by T. Seki (Osaka, Japan) and Y. Yanagihara and K. Noguchi (Kanagawa, Japan) . . . 245

Simultaneous quantitation of catecholamines and O-methylated metabolites in urine by isocratic ion-pairing high-performance liquid chromatography with amperometric detection
by Y.-P. M. Chan and T.-S. S. Siu (Pokfulam, Hong Kong) 251

High-performance liquid chromatographic analysis of josamycin in serum and urine
by M. Skinner and I. Kanfer (Grahamstown, South Africa) 261

High-performance liquid chromatographic method for the direct quantitation of oxy radicals in myocardium and blood by means of 1,3-dimethylthiourea and dimethyl sulfoxide
by P. S. Rao, N. Rujikarn and J. M. Luber, Jr. (New Hyde Park, NY, U.S.A.) 269

High-performance liquid chromatographic determination of indomethacin serum concentrations
by Y. L. Brown, R. J. Kandrotas, J. B. Douglas and P. Gal (Greensboro, NC, U.S.A.) . . 275

Measurement of the release of adenine nucleotides during platelet aggregation by small-bore-column isocratic high-performance liquid chromatography
by F. Bodola and C. R. Benedict (Galveston, TX, U.S.A.) 281

Rapid isolation of thymosin β_4 from thymosin fraction 5 by preparative high-performance liquid chromatography
by M. Badamchian (Washington, DC, U.S.A.), M. P. Strickler and M. J. Stone (Fairfax, VA, U.S.A.) and A. L. Goldstein (Washington, DC, U.S.A.) 291

Determination of noscapine and its metabolites in plasma by coupled-column liquid chromatography
by M. Johansson, A. Tufvesson Alm, H. Forsmo-Bruce and S. Jacobsson (Solna, Sweden) and D. Westerlund (Uppsala, Sweden) 301

Application of high-performance liquid chromatography to the analysis of propionyl-L-carnitine by a stereospecific enzyme assay
by A. Marzo, N. Monti and M. Ripamonti (Como, Italy) and E. Arrigoni Martelli (Rome, Italy) 313

Rapid and sensitive method for high-performance liquid chromatographic analysis of pterins in biological fluids
by I. Antonozzi, C. Carducci, L. Vestri, A. Pontecorvi and F. Moretti (Rome, Italy) . . . 319

High-performance liquid chromatographic evaluation of salicyloyl pyridixol and systemic metabolites in biological samples
by A. Marzo, A. Reiner, N. Monti and M. Ripamonti (Como, Italy) and C. Lucarelli (Rome, Italy) 325

Rapid identification of *Bacteroides* species by high-performance liquid chromatography
by L. Radin and A. Arzese (Genoa, Italy), C. Lucarelli (Rome, Italy) and G. A. Botta (Udine, Italy) 331

Improved high-performance liquid chromatographic determination of bacterial collagenase activity in ointments
by P. A. Biondi, F. Manca, A. Negri, C. Secchi and G. Tedeschi (Milan, Italy) and C. Lucarelli (Rome, Italy) 337

Improved high-performance liquid chromatographic analysis with double detection system for L-DOPA, its metabolites and carbidopa in plasma of Parkinsonian patients under L-DOPA therapy
by P. Betto, G. Ricciarello, M. Giambenedetti, C. Lucarelli, S. Ruggeri and F. Stocchi (Rome, Italy) 341

Determination of 5-hydroxytryptamine and 5-hydroxyindoleacetic acid in plasma by direct injection in coupled-column liquid chromatography with electrochemical detection
by B.-M. Eriksson and B.-A. Persson (Mölndal, Sweden) 351

Scale-up methodology for the preparative purification of peptide M
by M. Knight (Washington, DC, U.S.A.), M. P. Strickler and M. J. Stone (Fairfax, VA, U.S.A.), L. Chiodetti and S. Gluch (Washington, DC, U.S.A.) and T. Shinohara (Bethesda, MD, U.S.A.) 361

Preparative-scale high-performance liquid chromatography of omega-3 polyunsaturated fatty acid esters derived from fish oil
by J. M. Beebe, P. R. Brown and J. G. Turcotte (Kingston, RI, U.S.A.) 369

Author Index 379

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*)